

SPECIFICATION

Electronic Version 1.2.8

Stylesheet Version 1.0

[A METHOD FOR GRAPHICALLY DEPICTING DRUG ADVERSE EFFECT RISKS]

Background of Invention

- [0001] *Field of the Invention.* The present invention relates to an enhanced method for depicting the assessment and analysis of the risks of adverse effects resulting from the use of a particular drug, either alone or in combination with other drugs, nutrients, foodstuffs, beverages, toxins, chemicals, supplements, and other substances.
- [0002] *Description of the Related Art .* In September 1997, information regarding cardiopulmonary disease related to the use of fenfluramine and phentermine ("fen-phen") prompted the United States Food and Drug Administration (FDA) to request the manufacturers of these drugs to voluntarily withdraw both treatments for obesity from the market. Subsequent studies show a 25 percent incidence of heart valve disease apparently resulting from diet drug use. Thus, up to 1,250,000 people may have sustained heart valve damage from these diet drugs and the FDA indicates that this may be the largest adverse drug effect that the agency has ever dealt with.
- [0003] Current estimates are that some 2.2 million hospital patients had serious adverse drug reactions and more than 100,000 people die each year from adverse reactions to prescription drugs. Accordingly, federal officials have recommended that the FDA require hospitals to report all serious drug reactions to the agency. The Inspector General of the Department of Health and Human Services has also indicated that the FDA should also work to identify harmful effects of new drugs and encourage health-care providers to rapidly call the FDA with information about drug side effects. As new

drugs are introduced at increasing rates, the FDA will likely need additional resources to protect the public from hazardous drug side effects.

[0004] If one or two adverse drug reactions slip through the FDA's reporting process, the results can be tragic for some patients. That is especially true when the adverse reactions are rare but serious -- such as in the case of liver failure caused by medication. All drugs have the potential to harm or kill the people they are designed to help. An injection of penicillin can kill in minutes if the recipient is allergic to this life-saving drug. Even common aspirin can be deadly.

[0005] Clinical trials of a new drug often involve a few hundred patients and therefore may not reveal that a drug can cause serious injury or death in one patient in 10,000 or even 1,000 patients. Accordingly, it is critical for researchers and drug companies to be able to analyze and predict adverse reactions among patients in their studies.

[0006] In addition, in clinical trials for drugs used to treat diseases such as diabetes, which affects so many people and is difficult to treat, FDA officials often face tremendous pressure to accelerate their approval process. Often, in this "fast-track" process, cases of adverse drug effects may slip through reporting procedures.

[0007] An even bigger challenge to the FDA is the occurrence of adverse drug reactions after the drug is on the market. In this case the drug is prescribed to a much larger population of patients, many of whom are taking other substances such as extracts, nutrients, vitamins, hormones or drugs that might have an adverse effect with the prescribed drug.

[0008] Thus, there is a need for effective depiction of the risk analysis of adverse drug effects. Unfortunately, such a system has not been available.

[0009] U.S. Patent No. 6,120,443 to Cohen-Laroque, "Device For Determining The Depth of Anesthesia" discloses a method for determining the depth of anesthesia by acquiring a patient heart activity signal, converting the analog signal to a digital signal, sensing a periodic wave in the signal, determining the frequency of the periodic wave in the signal, determining the digital time interval series, calculating the fractal dimension of the digital series and calculating the depth of anesthesia based

on the fractal dimension. The reference also discloses a display means used to show the depth of anesthesia. The display means displays correlation dimension information.

[0010] U.S. Patent No. 6,108,635 to Herren et al., "Integrated Disease Information System" discloses a system that supports the development of new medical interventions for diseases. The system includes Explorer modules for discovering proposed interventions, designing clinical trials, performing pharmacoeconomic analysis, and illustrating disease progression for various patients over time including creating disease progression tutorials for a patient. The system also includes the ability to display correlations or relationships between selected input variables and disease progression measures. This graphical representation allows users to better understand and visualize patterns in the data that inform the clinical trial process.

[0011] U.S. Patent No. 6,151,581 to Kraftson et al., "System For And Method Of Collecting And Populating A Database With Physician/Patient Data For Processing To Improve Practice Quality And Healthcare Delivery" discloses a system and method for administering a patient management and health care management database. The database is built from data collected via patient surveys and contains data relevant to the clinical care of patients, the management of the practices to which the patients belong, and to the outcome of that health care and practice management. Once the data is collected from the patient, the data is entered in digital format and stored in a relational database. The data is then analyzed and prepared for dissemination. As part of the analysis, the data is scaled, scored and subjected to a battery of statistical tests that result in a series of correlative relations. The analyzed data is then presented in a printed or graphical format.

[0012] U.S. Patent No. 6,098,062 to Janssen, "Argument Structure Hierarchy System And Method For Facilitating Analysis And Decision-Making Processes" discloses a system and method for facilitating decision making. At the heart of the system and method is a computer program that links data representing argument structure units into a hierarchical argument structure. Each argument structure unit includes data representing a hypothesis and its corresponding counter-hypothesis, data

corresponding to grounds that provide a basis for the hypothesis or counter-hypothesis, data linking the grounds to the hypothesis or counter-hypothesis, and supporting data. Once all the inputs have been acquired, the hierarchical argument structure may be used for decision-making. Visualization programming code displays the support for and against each hypothesis at each node in the hierarchy.

[0013] U.S. Patent No. 6,076,083 to Baker, "Diagnostic System Utilizing A Bayesian Network Model Having Link Weights Updated Experimentally" discloses a diagnostic system, which utilizes a Bayesian network model having link weights updated experimentally. The system includes an algorithm for quantifying the strength of links in a Bayesian network, a method for reducing the amount of data needed to automatically update the probability matrices of the network on the basis of experiential knowledge, and methods and algorithms for automatically collecting knowledge from experience and automatically updating the Bayesian network with the collected knowledge. The reference also discusses structuring the display of evidence data so that it can be more easily filtered and accessed. In particular, the disclosure includes a graphical display means for displaying problems and diagnoses. The display means provides high level information with the ability to choose to display the underlying data that formed the basis of the high level information.

[0014] U.S. Patent No. 5,692,171 to Andres, "Method of Extracting Statistical Profiles, And Use Of The Statistics Created By The Method" discloses a method of extracting statistical profiles from a relational data base organized by relation, attribute, and index. The profiles are arranged in tables that are included in the database. The reference also discloses a method of profile extraction that can be adapted to any data type.

[0015] Japanese Patent No. 11282934, "Medicine Preparation and Medical Diagnosis Support Device" discloses a device that stores patient information, medicine information, examination information, etc. The device also includes a display processing part, which can display the medication information and examination data on the same time base.

[0016] Japanese Patent No. 10225500, "Drug Store and Drug Store Information System"

discloses a system to indicate the safety and effectiveness of two or more drugs prescribed at the same time. The system allows a user to select the drugs being prescribed from a medical supply table. The system then returns information about the drugs, which allows a doctor to judge their safety and effectiveness.

[0017] U.S. Patent No. 6,219,674 to Classen, "System for creating and managing proprietary product data" discloses systems and methods for creating and using product data to enhance the safety of a medical or non-medical product. The systems receive vast amounts of data regarding adverse events associated with a particular product and analyze the data in light of already known adverse events associated with the product. The system develops at least one proprietary database of newly discovered adverse event information and new uses for the product and may catalog adverse event information for a large number of population sub-groups. The system may also be programmed to incorporate the information into intellectual property and contract documents. Manufacturers can include the information in consumer product information that they provide to consumers or, in the case of certain medical products, prescribers of the medical products.

[0018] However, none of these references provides an enhanced method for depicting the assessment and analysis of the risks of adverse effects resulting from the use of a particular drug, either alone or in combination with other drugs, nutrients, foodstuffs, beverages, toxins, chemicals, supplements, and other substances.

[0019] Thus, there remains a need for an enhanced method for depicting the assessment and analysis of the risks of adverse effects resulting from the use of a particular substance, either alone or in combination with other drugs, nutrients, foodstuffs, beverages, toxins, chemicals, supplements, and other substances. There also remains a need for an enhanced method for depicting the assessment and analysis of the risks of adverse effects resulting from the use of a particular drug, either alone or in combination with other drugs, foodstuffs, beverages, toxins, chemicals, nutrients, supplements, and other substances.

Summary of Invention

[0020] The present invention pertains to an enhanced method for displaying assessment and analysis of the risks of adverse effects resulting from the use of at least one substance of interest, comprising: identifying the at least one substance of interest; selecting the profile of the at least one substance of interest related to the safety of the at least one substance of interest, using at least one filter; analyzing the risks of adverse effects resulting from the use of the at least one substance of interest using at least one data mining engine; and displaying the results of the analysis of risks of adverse effects resulting from the use of the at least one substance of interest in a format that permits perception of correlations. Such a format is preferably at least one format selected from the group consisting of a radar display for display of correlations, a sortable table, and a sortable line listing and where the format contains elements linked to data regarding the adverse effects.

[0021] In a preferred embodiment of the present invention, the enhanced method for displaying assessment and analysis of the risks of adverse effects resulting from the use of a drug of interest, either alone or in combination with other drugs, nutrients, foodstuffs, beverages, toxins, chemicals, supplements, and other substances comprises an input device whereby a user can identify the drug of interest, as well any other drugs, nutrients, foodstuffs, beverages, toxins, chemicals, supplements, and other substances; a selector for selecting the drug of interest's safety profile, adverse effect cases, drug reactions, and relationships therebetween, using at least one filter; at least one data mining engine preferably selected from the group consisting of (1) a proportional analysis engine to assess deviations in a set of the reactions of the drug of interest, (2) a comparator to measure the reactions of the drug of interest against a user-defined backdrop, and (3) a correlator to look for correlated signal characteristics in drug/reaction/demographic information; and an output device whereby a user can receive analytic results from the selector, and the at least one data mining engine, wherein the output device provides an enhanced depiction the assessment and analysis of the risks of adverse effects resulting from the use of a particular drug, either alone or in combination with other drugs, nutrients, foodstuffs, beverages, toxins, chemicals, supplements, and other substances.

[0022] This enhanced depiction of results of the assessment and analysis of the risk of

adverse effects resulting from the use of a drug of interest can be displayed using conceptual filters that can be modified based on pertinent variables, such as, age or outcome. Assessments of drug adverse effect risks are displayed in both a traditional tabular display format and a "radar-like" display format. The radar screen is populated with textual information and facilitates perception of correlations, sometimes permitting instant perception of correlations. The radar screen will allow zoom, peak thresholding, case report detail, drug/drug comparison and drug/therapeutic category comparisons.

[0023] The enhanced depiction method of the present invention relieves the user of the need to enter strings of criteria in testing a hypothesis regarding the assessment and analysis of the risks of adverse effects resulting from the use of one or more particular drugs. The present invention also relieves the user of the need to refer to off-line sources to test a hypothesis. Every dimension is afforded an intuitive visual display permitting a user to review and select one or more dimensions.

[0024] It will be appreciated that such a method for displaying the assessment and analysis of the risks of adverse effects resulting from the use of one or more particular drugs is advantageous to the various risk assessors who are tasked with making such determinations. Such risk assessors include governmental agents who perform such assessment for regulatory purposes, as well as agents of pharmaceutical manufacturers who are tasked with such assessments.

[0025] The present invention, which provides an enhanced method for displaying assessment and analysis of the risks of adverse effects resulting from the use of one or more particular drugs, offers an enhanced degree of analysis not previously available. This enhanced degree of analysis permits the identification of associations and, thus, potential causal elements regarding adverse effects resulting from the use of one or more particular drugs.

[0026] The present invention provides answers to several key questions that are essential to public health. For example, various safety groups, both government and private, are charged with monitoring the post-market behavior of drugs and determining "signals" that indicate a relationship among adverse reactions, demographics,

behavior, environmental exposure, dietary habits and other elements such as outcomes. Unexpected or previously unrecognized adverse drug effects can take the forms of single reactions, groups of reactions, or increases in a labeled reaction. Such adverse drug effects might be due to the higher exposure to the general population experienced in post-market therapy or such effects can be a reaction that has a demographic (genetic or otherwise) emphasis in an age or gender group.

[0027] Further, with efficient and effective analysis of adverse drug effects, pharmaceutical research and development professionals can learn more details of the reaction profiles of drugs and the at-risk populations who may be prescribed those drugs. This information would allow a more effective selection of lead compounds and would produce drugs with less risk of adverse effects.

[0028] The enhanced presentation of results in context-sensitive groupings (for example, reactions in a hierarchy or effects in a line) permits visual signal detection on the part of the user. The human brain is adept at perceiving certain features (for example, linear relationships, clustered data) that the present invention renders meaningful by linking to a logic framework. As an example, the present invention presents the correlation among variables of a set of adverse event cases, so that, if a high correlation is found, it can easily be determined which are the contributing elements. By providing a "temperature" scale for displaying a user-selected degree of correlation, the present invention allows the user to readily see the relative correlation.

[0029] Thus, the present invention allows for analysis of adverse drug effects with enhanced speed and flexibility. The present invention also offers new insights with regard to adverse drug effects and augments the existing processes of drug development.

[0030] Accordingly, it is an object of the present invention to provide an enhanced method for depicting the assessment and analysis of the risks of adverse effects resulting from the use of a particular substance, either alone or in combination with other drugs, nutrients, foodstuffs, beverages, toxins, chemicals, supplements, and other substances.

[0031] It is an object of the present invention to provide displays for assessments of drug adverse effect risks in both a traditional tabular display format and a "radar-like" display format. The radar screen is populated with textual information and facilitates perception of correlations, sometimes permitting instant perception of correlations. The radar screen will allow zoom, peak thresholding, case report detail, drug/drug comparison and drug/therapeutic category comparisons.

[0032] It is further an object of the present invention to provide hyperlinking of elements in the display so as to enable the user to drill down on those elements to obtain additional information. In a particularly preferred embodiment, the data presented in the display is hyperlinked to the specific details of the underlying cases.

[0033] Yet another object of the present invention is to provide a "radar-like" display which employs a visual scale in which brighter colors represent a higher degree of correlation and duller colors represent lesser degrees of correlation. The method of the present invention further allows the user to control the display of this "temperature-like" display. This user-control feature permits a user to establish threshold values for the display of mean and range values, thus, facilitating perception of correlations.

[0034] It is another object of the present invention to provide a method for linking statistics or associations with the underlying supporting cases.

[0035] It is yet another object of the present invention to provide a method of presenting data in a way to permit coherent visual processing.

[0036] Further, it is another object of the present invention to provide filtering devices in great numbers without requiring extensive keyboard entry or outside references.

[0037] It is still another object of the present invention to provide a method for a user to dynamically change scales to uncover the threshold range of difficult-to-predict correlations.

[0038] A greater understanding of the present invention and its concomitant advantages will be obtained by referring to the following figures and detailed description provided

below.

Brief Description of Drawings

- [0039] Figure 1 is chart indicating the page flow of the present invention;
- [0040] Figure 2 is an overview of the present invention;
- [0041] Figure 3 is a depiction of a home page of the present invention;
- [0042] Figure 4 is a representation of a filter selection page of the present invention;
- [0043] Figure 5 is a representation of a selector page of the present invention;
- [0044] Figure 6 is an illustration of an exemplary pedigree screen of the present invention;
- [0045] Figure 7 is a depiction of an exemplary reactions table in the profiler component of the present invention;
- [0046] Figure 8 is a representation of a concomitant drugs table in the profiler component of the present invention;
- [0047] Figure 9 is a depiction of a demographics table in the profiler component of the present invention;
- [0048] Figure 10 is an illustration of a report dates table in the profiler component of the present invention;
- [0049] Figure 11 is a representation of an outcomes table in the profiler component of the present invention;
- [0050] Figure 12 is a depiction of a reaction filter screen of the present invention;
- [0051] Figure 13 is an illustration of a correlation results screen of the present invention;
- [0052] Figure 14 is a representation of a correlation details screen of the present invention;
- [0053] Figure 15 is a depiction of a case details screen of the present invention;

05684586 050201
T02050 030501

[0054] Figure 16 is an exemplary illustration of a radar screen display of the present invention;

[0055] Figure 17 is an illustration of a proportional analysis selection screen of the present invention;

[0056] Figure 18 is a representation of a proportional analysis results screen of the present invention;

[0057] Figure 19 is a depiction of a tabular version of a proportional analysis screen of the present invention;

[0058] Figure 20 is an illustration of a comparator screen of the present invention; and

[0059] Figure 21 is a representation of a case list of the present invention.

Detailed Description

[0060] The present invention provides an enhanced method for displaying the assessment and analysis of the risks of adverse effects resulting from the use of one or more particular drugs, either alone or in combination with other drugs, nutrients, supplements, vitamins, foods, beverages, and other substances.

[0061] The enhanced method for displaying the assessment and analysis of the risks of adverse effects resulting from the use of one or more particular drugs preferably includes, in one embodiment, radar-like displays incorporating intuitive displays arranged in a user-selectable configuration. For example, in one preferred embodiment of the present invention, the radar-like display can employ a visual scale in which brighter colors represent a higher degree of correlation and duller colors represent lesser degrees of correlation. The method of the present invention further allows the user to control the display of this "temperature-like" display. This user-control feature permits a user to establish threshold values for the display of mean and range values, thus, facilitating perception of correlations.

[0062] In another preferred embodiment of the present invention, the enhanced method for displaying the assessment and analysis of the risks of adverse effects resulting

from the use of one or more particular drugs preferably includes the representation of tree structures and pathways applied to the various dimensions of the adverse effect risk assessment.

[0063] For purposes of the present invention, a substance shall be understood to include one or more elements of the set of drugs, foodstuffs, beverages, nutrients, vitamins, toxins, chemicals, hormones, and supplements.

[0064] Another preferred embodiment of the enhanced method for displaying the assessment and analysis of the risks of adverse effects resulting from the use of one or more particular drugs preferably includes context-sensitive pull-down menus to view and select filters.

[0065] Still further, another preferred embodiment of the present enhanced method for displaying the assessment and analysis of the risks of adverse effects resulting from the use of one or more particular drugs preferably includes hyperlinking of elements in the display so as to enable the user to drill down on those elements to obtain additional information. In a particularly preferred embodiment, the data presented in the display is hyperlinked to the specific details of the underlying cases.

[0066] Another preferred embodiment of the enhanced method for displaying the assessment and analysis of the risks of adverse effects resulting from the use of one or more particular drugs preferably includes the ability to group data in pre-defined clusters in both tables and radar screens so as to enhance a user's pattern recognition and visual integration.

[0067] The primary components of a preferred embodiment of the present enhanced method for displaying assessment and analysis of the risks of adverse effects are a combination of the following: one or more integrated databases; a selector for selecting at least one drug for analysis (based on the generic, brand names or therapeutic category); a profiler for displaying statistics that describe behavior for the drug in multiple dimensions; a series of at least two filters and the means to control the at least two filters individually and in combination; at least one of three or more data mining engines selected from the group of a correlator, a proportional analysis

09691596-050201

or filter in various dimensions of the risk assessment at the Filter in Various Dimensions Box 121.

[0073] If the user has chosen to access more details, then the user is preferably presented with multiple dimensions of the risk assessment from which to access more information. Preferred dimensions of risk assessment include, but are not limited to, Reactions More Details Box 115, Concomitant Drugs More Details Box 116, Demographics More Details Box 118, Report Dates More Details Box 118, and Outcomes More Details Box 120.

[0074] Preferably, if the user has chosen to filter in various dimensions of risk assessment, then the user is preferably presented with multiple filters for the dimensions of the risk assessment. Preferred filters of dimensions of risk assessment include, but are not limited to, Reactions Filters Box 121, Concomitant Drugs Filters Box 122, Demographics Filters Box 123, Report Dates Filters Box 125, and Outcomes Filters Box 126.

[0075] The preferred components of the present invention are illustrated in Figure 2, which provides an overview of the method of the present invention. The user preferably accesses the system of the present invention by means of Home Page 200. From Home Page 200 the user can proceed to the Selector 201, where the user can select a drug for analysis. Having selected the drug of interest, the user can then preferably proceed to the Profiler 202, which preferably displays statistics that describe the behavior of the drug of interest. From the Profiler 202 the user can then preferably proceed to employ one or more Filters 203, which permit recalculation of the statistics by selecting among the available variables. Once a set of cases is determined, for example, by the use of one or more filters, the cases can then preferably be submitted to at least of three or more Data Mining Engines 204. The output from the data mining engines is then preferably displayed in a Viewer 205, which can present the data in a variety of formats, including, but not limited to a sortable table, a sortable line listing, and a radar screen, thus, allowing rapid identification of signals and providing the user the ability to drill down to individual case details.

102050-05001

- [0076] Alternatively, in another preferred embodiment, from Home Page 200 the user can choose a profile from the Profiler 202, apply one or more Filters 203, process the set of cases using the Data Mining Engines 204 and display the results with the Viewer 205.
- [0077] Preferably, the present invention operates on at least one of two integrated databases: an external public database characterized by breadth of data across all drugs and a database containing internal data of an organization or an individual characterized by increased detail with regard to one or more specific groups of drugs. It will be appreciated that a database containing the internal data of an individual can refer to a number of different situations, including but not limited to the biological/medical/genetic/environmental exposure/behavior/demographics/drug sensitivity of an individual.
- [0078] In both cases, the source and purpose of the data may vary, including post-marketing surveillance, clinical trial data, health care system data, research databases, and literature, among others.
- [0079] The various aspects of the enhanced method for displaying the assessment and analysis of the risks of adverse effects resulting from the use of one or more particular drugs, for example, the radar-like screens of the correlator and the proportional analysis engine, and the sortable line listings of cases, permit a user to undertake a logical drilldown in a risk assessment so as to view additional and more specific data, while maintaining traceability through the multitude of cases.
- [0080] The public database preferably is at least one database selected from either a combination of one or more of the FDA's Spontaneous Reporting System (SRS)(after to November 1997)) and the FDA's Adverse Event Reporting System (AERS)(after to November 1997)), the World Health Organization adverse event database, or other country-specific regulatory or epidemiological databases, such as the UK Advert system and the General Practice Research Database (GPRD). These public databases are updated regularly as they release new case data. In a related invention, which is a particularly preferred embodiment of the present invention, the present enhanced method for displaying assessment and analysis of the risks of adverse effects relies

upon a derivative of these public databases that has cleaned, parsed in to a relational database, and mapped to known dictionaries, and standardized for efficient searching and query defining. This preferred derivative database has over 2 million cases representing 30 years of adverse events as reported to regulatory authorities. Additionally, the derivative database links the adverse event (AE) case data to Medical Dictionary for Regulatory Activities (MedDRA), Coding Symbols for a Thesaurus of Adverse Reaction Terms (COSTART), and World Health Organization (WHO) Adverse Drug Reaction Terminology (WHOART), among others for reactions, and the National Drug Code Directory (NDCD), Orange Book or WHO Drug dictionaries for drugs. The invention includes the facility to substitute and manage standard dictionaries for all dimensions.

[0081] In a preferred embodiment, the present method also operates on a database containing internal data of an organization. For example, such an internal database could be the proprietary database of a pharmaceutical company or the contemporaneous database of a clinical investigator during the course of clinical trials upon a drug.

[0082] It will be appreciated that one preferred embodiment of the present invention utilizes a log-on screen. Preferably, access to the present invention is provided by means of the Web. The present method preferably supports all browsers including Netscape and Internet Explorer for access. In a particularly preferred embodiment, different URLs are used for the public database and for the internal database. This allows operating in two databases concurrently if two instances of the Web browser are opened. It also allows virtually unlimited simultaneous processes, and simultaneous processing at various locations.

[0083] The use of multiple sessions also enable a range of comparisons, in each and every dimension. The "differencing engine" or comparator provides immediate information on similarities and differences.

[0084] In another preferred embodiment of the present invention, a home screen is used to launch searches, and review results of the analytical engines and prior work. For example, from the home screen a user can (1) select a drug to study by either name or

09681536-050001

by therapeutic category, (2) recall a previously saved filter (that was created, named and saved previously, (3) review previously submitted analyses, or (4) invoke certain data mining engines directly. An exemplary version of a home page of the present invention is provided in Figure 3. From this home screen, the user can use field 301 to select a drug to study by generic name, trade name, or therapeutic category. The user can also use field 302 to recall a previously saved query (called a filter). Further, the user can use field 303 to recall previously submitted analyses. Additionally, the user can use field 304 to invoke the proportional analysis engine.

[0085] The home page is preferably the user's command center for analysis. The home page is preferably always accessible from any other screen.

[0086] Preferably, the home screen has four areas. The first area is a link to the selector, thus, allowing a user to easily reach the drug selection screen through any level of detail on a drug. The second is a filter area. The user can view and apply previously saved filters. The third section is the data mining engine section which allows a user to invoke one or more of the data mining engines. The fourth area permits the user to review previously generated analyses.

[0087] With regard to the selection of a drug, this feature allows a user to select at least one drug to study and to search for information on that at least one drug by using either its generic name, its trade name, its therapeutic category, or its chemical name. In addition, the invention provides the ability to develop specific other valuable taxonomies, such as a "super generic" including all salts of a drug, or a sub-brand, for example distinguishing between a once a day version of a drug from a once a week version of the drug.

[0088] Concerning selecting a previously saved query, this feature (referred to as a filter) is a preferred paradigm to reduce the routine of inputting a previously employed and saved query. By establishing parameters for searching, a user does not need to define ad hoc queries. Knowledge of the pharmacovigilance domain is used to present users with filter/query-building interfaces that are more in line with the thought processes and paradigms employed by such users. A user can preserve the set of parameters of a query (a filter) each time he/she refines a profile, and further, a user can employ a

09681586-050204

filter he/she has developed in a previous search the next time he/she wishes to view the same or an updated set of cases.

[0089] For example, Figure 4 provides a representation of a preferred filter screen. Various preferred fields of a filter screen are presented, including, but not limited to, Reactions (field 400), Concomitant Drugs (field 401), Demographics (field 402), Report Dates (field 403), and Outcomes (field 404).

[0090] In invoking a saved filter, a user is offered the option of viewing or applying (querying with) a saved filter, and a pull-down menu allows a user to select one of the filters previously created and saved. Pushing the "View" button allows a review of the specific details of the filter. In the example provided, a user had created and saved a filter he/she had labeled "Filter 2" for a search on Candesartan Cilexetil. The search results show the drug's Reactions, including its MedDRA Hierarchy Group (System-Organ-Class (SOCs), etc.), and a pull-down menu showing the specific reactions (ear and labyrinth disorders, for example) included in the filter.

[0091] Case sets, as well as drug sets, can be created, named, and saved similar to filters. Because these case sets generate a list rather than a logic description, viewing and changing are performed with a list manager. Filters, drug sets, and case sets can all be combined or merged to provide a rich set of functions, and great flexibility.

[0092] The preferred parameters of a filter include reactions (listed as "included" or "off"); concomitant drugs (listed or "off"); demographics (listed as per previously set brackets or "off"); report dates (listed or "off"); and outcomes (listed or "off"). If a user wishes to apply this saved filter as his/her current query, he/she would click on the "Apply" button. At this point, a user would be taken to the profile screens for that drug and that set of filters.

[0093] With regard to the review of the previous analysis aspect of the present invention, this section of the home page provides information on previous analyses a user has run using the correlator engine of the present invention. As illustrated in Figure 3, the correlator engine notices provide information on analyses that have been previously completed including date and time, task number, and generic drug. Each listing ends

0961586-05001
T02300-935T3960

with a hyperlink that a user can employ to view the results of the search. A "delete" function is preferably provided to manage this list.

[0094] Concerning the proportional analyzer aspect of the present invention, this component looks for large or small deviations in the reactions counts for a set of drugs, i.e., comparing drugs to those in their own therapeutic category or to all drugs. With a preferred embodiment of the present invention, a user has an option to compute for a therapeutic category using a pull-down menu. A user also has an option of selecting Bayesian filtering. Bayesian filtering employs a statistical cut-off threshold to reduce the affect of rows or columns with a very low number of cases. That is, drugs or reactions accounting for less than a certain percent of cases or fewer than a set number will be deleted from the matrix (and so noted on the results screen).

[0095] A user preferably has two options in running this analysis: (1) he/she can compute information for each drug's reactions in comparison with all of the drugs in the system or (2) a user can run an analysis by comparing the selected drug's reactions only with those of other drugs in the same therapeutic category. Results are preferably presented concurrently on a separate screen.

[0096] An additional preferred aspect of this home page is a comparator, which is available when a user is accessing optionally provided clinical trial data from a drug label, or from the clinical trial data of an internal database. The comparator compares potential and actual adverse effects of drugs in the pre- and post-market environments.

[0097] The preferred home page of the present invention also provides a user with the options to add a user, manage preferences, manage the group of inserts, and to log out, among others.

[0098] If a user has selected a query at the Home page, he/she will initiate a query for a drug using the drug's generic name, its trade name, its therapeutic category, its chemical name, or other custom-defined categories. The search invokes the selector page of the present invention. A user selects a drug by clicking on the generic drug

0963136-05001

link which then takes a user to the profile and general statistics regarding the selected drug. A user starts his/her search on the home screen, and then continues it on the selector page, by entering or selecting the category of drug he/she wants to search: the generic name, the trade name, the therapeutic category or the custom-defined categories. The therapeutic category field preferably has a pull-down menu to help identify and select the desired field.

[0099] An exemplary Query Screen page is illustrated in Figure 5, where a user has decided to search the therapeutic category of angiotensin converting enzyme (ACE) inhibitors, as defined by the drug dictionaries. (Note: In this case the FDA taxonomy places certain drugs known as ATII drugs in the ACE category.) Here, the user has chosen not to use the generic name field 500 or the trade name field 501, but rather has chosen the therapeutic category field 503. The present system returns with the hits corresponding to the selected therapeutic category and are displayed in the query screen. In this example, 22 drugs matching the search criteria were found in the "ACE Inhibitors" category. The drugs are listed in alphabetical order by their generic name. For each generic drug on the list, all trade names and all relevant therapeutic categories are presented in pull-down menus. Optionally, custom-defined categories can also be shown. The search results also allow access to the drug's "pedigree," or lexical mapping information, indicated by a question mark link.

[0100] Preferably, a user can stop browsing drugs and go directly to the profile by selecting and applying a previously stored filter.

[0101] With regard to the pedigree function, if a user selects the pedigree icon for the selected drug (the question mark in this example), a user is presented with the drug's pedigree, which shows the way the drug has been mapped in a drug dictionary and thesaurus.

[0102] An exemplary pedigree screen is presented in Figure 6. This exemplary pedigree screen provides a number of preferred fields indicating the cataloging of the data in the system of the present invention. Preferred fields include, but are not limited to, Map To (field 600), Verbatim (field 601), Source (field 602), Incidents (field 603), Case Count (field 604), QEDRx Processing (field 605), Cross-Reference (field 606), and

[0105] The profiler aspect of the present invention permits a user to navigate various dimensions of the selected drug's safety profile and view cases, concomitant drugs, reactions, demographics, outcomes, and time intervals using specified filters. Once a user is satisfied with the cases profiled, the set of cases satisfying the filter criteria can then be submitted to the various data mining engines, including the Correlator Engine (CE), Proportioning Engine (PE) and Differencing Engine (DE). Each data mining engine is provided with a set-up and a verification step (by means of a page set of input parameters). For example, the CE may further weight the different dimensions.

[0106] It will also be appreciated that the profiler of this invention allows for continuous adjustment and addition to the dimensions. For example, a preferred embodiment includes "Repeat Source". Others may contain laboratory results. The invention permits expanding and contracting both the profiler and the at least two filters as the data changes.

[0107] As noted above, in the selector component of the present invention, each of the drugs in the generic name category is preferably presented in a format that indicates a hyperlink. Clicking on a generic drug (in the previous example Candesartan Cilexetil), the multi-dimension profile screen is invoked by clicking on a generic drug (in the previous example Candesartan Cilexetil).

[0108] The idea of profiling a drug is complex, because of the multiple dimensions. The invention's profiler separates presenting data on the selected drug into several different categories and preferably "billboards" the top ten for immediate visibility. It will be appreciated that the user can specify any number for "billboarding." At the top of the screen are the generic name of the drug (preferably with a hyperlink to its pedigree), all the trade names associated with the drug, and all of the therapeutic categories to which it belongs.

[0109] The profile feature of the present invention is used to display statistics that describe the effects of the drug in multiple dimensions. Each set of data is preferably presented in a separate table, headed by an index tab. The preferred data sets include, but are not limited to: (1) Reactions; (2) Concomitant Drugs; (3) Demographics; (4) Report Dates (for example, dates logged by FDA as report dates for

096156-036156

SRS and AERS); and (5) Outcomes.

[0110] For each dimension there are key actions: all allow filtering and delving for more details. The filter action allows a user to set and activate filters for that dimension. The "More Details" action brings up all the values that have appeared only in the top 10 billboard style on the main page.

[0111] For certain dimensions, for example reactions, the hierarchy of the dimension can be selected to change the billboard and detailed views. In the case of reactions, MedDRA contains a five level hierarchy. Other dictionaries use two to four levels. The present invention accommodates the full range of hierarchies.

[0112] Preferably, the profiler feature of the present invention allows grouping concomitant drugs by therapeutic category, chemical class, or other custom-defined class.

[0113] With regard to the Reactions dimension, the profiler component of the present invention preferably shows reactions to the drug that is being queried. This dimension refers to suspected adverse reactions to the selected drug that were reported. A suitable reactions table is provided in Figure 7. In this figure, to the right of the Reactions tab is a pull-down menu labeled "View" 700, followed by a filter hyperlink 701. By utilizing the pull down menu, a user can choose among multiple different levels of MedDRA. Of these multiple different levels of MedDRA, four are particularly preferred. These are System, Organ, Class (SOC), High Level Group Term (HGLT), High Level Term (HLT), and Preferred Term (PT).

[0114] In Figure 7, a user has chosen the HLT option. The window in the pull-down menu indicates that there are 256 HLTs out of the total of 1495 HLTs in the current version of MedDRA. The Reactions Table 702 shows the Top 10 HLTs of the 256. In this case, the reactions include hypertension, disturbances in consciousness, and so forth. For each of the reactions, the table presents the Reaction Count (the number of times this reaction was listed in the database) and the percentage of reactions that this number constituted in the set of reactions for this drug, based on incidents of reactions (not cases).

102050 "0501
0901396
0901396

[0115] At the bottom of the Reaction Count and % of Reactions columns are numbers showing the number of incidents of the reactions at the Top 10 HLTs (488) and the Total Reactions across all of the 256 HLTs (in this case, 1752), 703 and 704, respectively.

[0116] The ability to browse statistics, up and down a hierarchy, and within real time, is important to keeping risk assessment hypothesis setting and testing within a short period of time. The invention provides extensive associative tables and reverse indexing to enable such rapid analysis.

[0117] A hyperlink offering more details concurrently follows the Reactions table, and brings up a separate page with all details of this dimension.

[0118] It will be appreciated that since Reactions, Concomitant Drugs, and Outcomes are summarized at the event level, the resultant collection of cases will be different if more than one event is associated with a single case. For example, if two reactions are recorded in a single case, and both of those reactions parent to the same MedDRA SOC, then they will account for two events, and yet would yield only a single case in the case listing.

[0119] In a preferred embodiment of the present invention, case level percentages and percentage relative to drug exposure are also available in the profiler component of the present invention. With regard to the Concomitant Drugs dimension of the profiler of the present invention, this dimension describes drugs that were also prescribed in the cases in which the target drug was found. A suitable example of a Concomitant Drugs table is provided in Figure 8.

[0120] In this figure, the Concomitant Drugs Table 800 lists the top 10 drugs in the concomitant category. In this example, hydrochlorothiazide, aspirin, and furosemide were among the drugs found in combination with Candesartan Cilexetil in the adverse reactions reported to the FDA.

[0121] The table divides the cases of concomitant drugs into two groups: Suspect and Non-suspect (fields 801 and 802 respectively). When an adverse reaction report is filed, certain drugs in the case may be indicated as suspect. When considering

of reports for the drug Candesartan Cilexetil that fall within that time interval. By selecting the more details link, a user can obtain the breakdown of the reports by individual years.

[0127] In the Outcomes dimension of the profiler of the present method, case outcomes are listed. An appropriate outcomes table is presented in Figure 11. Preferred categories include serious outcomes such as congenital anomaly, death, and disability, as well as other outcomes. Serious outcomes are preferably presented in red, while less- or non-serious outcomes are in black. The Outcomes Table provides a table of outcomes (field 1100), a count (field 1101) and percentages of the outcomes (field 1102) in each category, as well as totals of serious and non-serious outcomes.

[0128] The filtering feature of the present invention is a paradigm that reduces the routine of constructing ad hoc queries. This filtering feature is context-sensitive and relieves a user of the burden of repeatedly defining the parameters of the queries. Filtering allows a user to formulate queries in a way more consistent with paradigms used by medical professionals, selecting among the active cases and using standard dictionaries such as MedDRA and National Drug Code Directory. This filtering features preferably allows a user to apply and view filters individually, set filters as a group and apply globally, or save and apply filters at a later time.

[0129] Data is compiled by the filters selected for each analysis. In the above example, filters were established for the reaction query. One of the screens in the profiler component was the reactions dimension, providing the Top 10 SOC's for the drug Candesartan Cilexetil. At the top of the table was a pull-down menu with "View" selected, also provided with a filter hyperlink.

[0130] Each of the data sets in the Profiler (Reactions, Concomitant Drugs, Demographics, Report Dates, and Outcomes) provides a user with the opportunity to establish filter parameters in any order. In a preferred implementation, the invention tabs the individual filters for convenience, and allows merging with other filters.

[0131] An exemplary filter applied as to reactions is provided in Figure 12. This figure provides the list of Reaction Filters available for profiling. The filter is based on the

MedDRA hierarchy and begins at the SOC level.

[0132] The mechanics for working with filters is common to all dimensions. A user may click on any or all of the reactions they would like to have included in the filtered reaction profile.

[0133] In the example of Reaction filtering, clicking on an SOC brings up the HLGTs for that SOC and allows selection at that level.

[0134] In a preferred embodiment filtering can be done at all levels.

[0135] It will be appreciated that for the more complex filters, such as the reaction filter, a range of user friendly aids is provided. For displayed MedDRA leads, preferably a tree is used. When it is collapsed, an open box preferably means no selections lower in the hierarchy have been identified, a check means all lower selections in the hierarchy have been identified, and a new query box is used to indicate unchecked box(es) somewhere below in the hierarchy.

[0136] Another preferred feature of the present invention is content-based pre-filters. To make it easier to switch-off indication-related adverse drug reactions, an "indications-related" button is preferably provided in the selection. For labeled adverse effects, of which there could be hundreds, the invention preferably provides tables (in this case with data from drug labels) to switch off all of the labeled reactions. This quickly focuses the user's attention on "unexpected" reactions.

[0137] Preferably on the profiler component, the present invention monitors the contents of each filter as it is built. At any point, the filter can be saved as an entirely new filter or by overwriting an old one, or changing and saving an incremental filter. This permits fine tuning of hypotheses regarding adverse drug reactions.

[0138] The filter for the concomitant drugs dimension allows selecting or deselecting any and each of the concomitant drugs reported in the profiled set of cases. Similar to reaction filtering, the concomitant drug dimension filter preferably provides a context selector (for example, to switch out a whole therapeutic category).

[0139] The demographics filter allows selections of generational or individual age

brackets, and male/female selections as well. Generational filters are preferably user definable.

[0140] The report dates dimension allows selection by bracketed years. In addition, in another embodiment of the invention, the report dates filter incorporate a link to a drug's birth date and allow filtering by "first six months," "first two years," etc. A table of drug birth dates relieves the user of the need to separately enter those dates.

[0141] The Outcome filter allows individual outcome selection, or by serious/non-serious grouping. For internal database adverse events, if a custom seriousness set is defined, this dimension will be user definable.

[0142] The analysis provided by the method of the present invention finds "signals" such as anomalies in a random population, a change against a known background, or a coherent target in a noise background. This is accomplished by at least one of three or more data mining engines: the proportional analysis engine (PE), the comparator (differencing engine or DE), and the correlator. In a preferred embodiment, the proportional analysis engine can be invoked from the home screen, as can be the comparator, for selected data. The correlator is invoked after filtering cases from the profile page.

[0143] The correlator looks for the association of characteristics in literally millions of pieces of drug/reaction/demographic information concurrently.

[0144] Too often in risk assessment, important correlations are hidden by surrounding background "noise" that obscures connections among data elements. Using a multidimensional vector analysis, the correlator measures the degree of association among pairs of values (for example, a drug and a reaction, an age and an outcome, etc.). The correlation algorithm is user selectable and definable. The preferred version uses a Pearson product-moment correlation known conventionally as " R^2 ". Other algorithms can also be used. The invention preferably applies the correlation after filtering, greatly enhancing the signal and reducing noise.

[0145] As a preferred example, the profiler screen can provide a number of hyperlinks choices, including "Apply Filter" and "Compute Correlations." Selecting "Compute

102050" 925T8950

[0148] In addition to viewing the table listing online, a user may also preferably select to review the results using a "radar-screen" correlation viewer. On the correlation screen, after the "Below are the top 200 correlated terms for your analysis . . .," there is preferably a hyperlink that provides the option of viewing the results with the correlation viewer. In addition to viewing with the correlation viewer, a user is also preferably presented with options to save the file.

[0149] Two other information screens preferably provide additional information provided by the correlation engine. From the correlated terms screen, a user is preferably presented with hyperlinks comprising all of the numbers in the Rank column. A significance (to a user-selectable "P" value) is also preferably provided. These hyperlinks provide a link to individual case lists. An exemplary correlation details screen is provided in Figure 14.

[0150] The Correlation Details screen of Figure 14 provides the data for each of the cases included in that pair of correlated terms. For example, if the term pair in the Correlated Terms Screen was "Female" and "Candesartan Cilexetil," this screen provides the pertinent information for all of the cases where those two terms were paired. In this example, there were 18 cases where renal function analyses were correlated with Candesartan Cilexetil. For each case, preferably the following information is provided: the case ID (field 1401); the gender of the patient (field 1402); the Manufacturer's Control Code (field 1403); the FDA Report Receipt Date (field 1404); the patient's age (field 1405); the other drugs the patient was taking at the time of the incident(s) (field 1406); the patient's reaction(s) to the medications (field 1407); and whether the outcome was Serious (yes or no) (field 1408). By selecting these cases, the user can then profile the set of cases.

[0151] Additionally, to learn the details of a specific case, a user preferably can click on the case ID number of any case on the Correlation Details screen. The resultant information is preferably presented in a case details screen. An suitable case details screen is presented in Figure 15.

[0152] The Case Details screen of Figure 15 provides detailed information on each specific case. In addition to standard information such as the patient's case ID (field

1501), gender (field 1502), and age (field 1503), it preferably includes Reactions (field 1504)(including detailed information in the As Reported, Preferred Term, High Level Term, and High Level Group Term categories); Concomitant Drugs (field 1505)(each listed by Name, Dose, Route, and Suspect Status); Outcomes (field 1506); Manufacturer Control Code (field 1507); Manufacturer Date (field 1508); Adverse Event Date (field 1509); Report Type (field 1510); Report Source (field 1511); Case Source (field 1512), and Narrative (field 1513), if any. All data, including lab test and genetic information can be encoded and displayed.

[0153] It will be appreciated that the above-identified information is not the only information that can be provided; extra information fields may be also provided.

[0154] The adverse effect analysis result of the present invention are preferably presented in a format that provides both traditional tabular displays (line listings) and innovative "radar-like" displays. By populating a radar screen with textual information, a user moves from the cumbersome reading of printouts to the instant perception of correlations directly on the screen. Once a signal is identified, a case browser permit a user to move through user-defined sorting to the key cases involved. Once again the synergistic aspects of the invention come into play. A "Therapeutic Category" or "Labeled Reaction" selector can group the data on the radar screen to enhance the signal. An exemplary radar screen display is presented in Figure 16.

[0155] The proportional analyzer engine of the present invention monitors outliers among reactions for drugs, for example, by comparing drugs to all drugs or those in a therapeutic class. The proportional analyzer engine can employ a variety of algorithms, including, but not limited to, PRR, ODDS ratio, and PRE, among others.

[0156] The proportional analyzer is preferably invoked from the home screen. A user is, in a preferred embodiment, prompted to select a therapeutic category for analysis by the proportional analyzer engine. Alternatively, a drug or a drug set can be selected. A user can select the therapeutic category that contains the drug he/she wishes to analyze. Bayesian filtering is preferably available as an option to remove noisy results due to lower case counts from the analysis.

All pairs of the dimensions of the analytical engine (for example, reaction and outcomes) can be analyzed. Even within the cases of a single drug, the reactions and concomitant drugs could be proportionally analyzed.

[0162] In addition to the graphic display, the proportional analyzer also shows these data in a tabular form. Figure 19 is the tabular presentation of the proportional analysis results. In this table, the location of the drug (field 1901) and its reaction (field 1900) in the matrix are indicated by numbers for row and column, row indicating the reaction and column signifying the drug of interest. The remaining three columns in the table preferably indicate the reaction count (field 1902)(with a hyperlink to the cases themselves), the expected reaction count (field 1903), and the Relative Ratio (field 1904). The entries are ranked in descending order, with the highest ratios listed first. The columns can preferably be sorted by clicking on their headings.

[0163] As in all tables, from the selector to the correlator, numbers are hyperlinked to the case-list. In the proportional analysis engine, all HLTs are available.

[0164] Such an enhanced display of results of the assessment and analysis of the risk of adverse effects resulting from the use of a drug of interest permits a user to reach rapid conclusions by enhancing signal processing of the results by the user. In the display of the present invention, a user can look across all data visualizations and quickly discern patterns which might otherwise have been obscured beneath a large amount of data.

[0165] The enhanced display of the assessment and analysis of the risk of adverse effects resulting from the use of a drug of interest of the present invention exploits a particular strength of the human mind, pattern recognition. The enhanced display of the present invention provides a user with data in a format that permits rapid pattern recognition. For example, when a user is assessing causative elements, the user must first determine whether a particular parameter is significant. The preferred format of the display method of the present invention permits a user to rapidly ascertain the significance of a correlation. For example, the correlator can be set to display a bright dot for a user-defined level of correlation. The graphical user interface of the present invention will then display the row of all of the primary putative causative elements. If

there are other darker dots in that row, there might be another causative element in the corresponding column. In this example, if a bright dot is under the primary drug and a darker dot is under one age group, that would suggest that the drug might be demonstrating an adverse effect with regard to that age group.

[0166] Another significant aspect of the enhanced display of the assessment and analysis of the risk of adverse effects resulting from the use of a drug of interest of the present invention is that the visualization data is further linked to additional information. Conventionally, a graphical user interface has been used to display information, but not to facilitate decisions. The present invention seeks to provide not only information, but suggest approaches for hypotheses and decisions. For example, with a graphic display method of the present invention, a bright dot indicates a high correlation; that bright dot of the graphical display of the present invention is linked to the putative causative elements, to the cases, to the profile, and to the filter.

[0167] The graphical display of the present invention also preferably provides multiple hyperlinks to information and permits toggling to a line listing format for the information. This format permits a user, by means of multiple sorting in a columnar line listing, to drill down into a data set.

[0168] Such sorting permits a user to utilize hierarchical processing with regard to the various parameters involved in the sorting. Upon completion of the hierarchical processing, the user can then exploit the hyperlinking which preferably underlies each datum.

[0169] Thus, in a preferred embodiment, the graphical display of the present invention offers a user a visual display for pattern recognition; a line listing correlated with the visual display; dynamic sorting of line listing; and hyperlinking of the elements provided in either the visual display or the line listing.

[0170] The comparator or differencing engine screen in the preferred offering offers three sets of analyzed data: Pre/Post Market data, Other Post-Market Reaction, and Other Clinical Trial Reaction. An exemplary comparator screen is provided in Figure 20. The Pre/Post Market data is preferably organized into a series of columns in a first

table (field 2000), providing the information, including Reaction HLT (field 2001); Clinical Trial Reaction (field 2002); Clinical Trial Percentage (field 2003), Clinical Trial Adjusted Percentage (field 2004); Post Market Reaction (field 2005); Post Market Percentage (field 2006); Post Market Adjusted Percentage (field 2007); and Difference Ratio (field 2008). The adjusted percentages account for proportions of those reactions that are common in both pre- and post-market reporting. The second table (field 2009) lists Other Post-Market Reaction (field 2010) and each reaction's Post-Market Percentage (field 2011). This information represents data available in the integrated public database. The third table (field 2012) provides Other Clinical Trial Reaction (field 2013) and each reaction's Clinical Trial Percentage (field 2014). This information indicates whether this reaction was mentioned on the manufacturer's package insert.

[0171] The comparator engine of the present invention is a differencing engine that is applied to measuring one drug's reactions, both pre- and post-market. This engine is essentially a "proportion of proportions" and is preferably limited to situations where: labeled adverse effect data can be quantified, terms can be mapped to MedDRA, and a useful number of reports are available for reactions, both pre- and post-market. The comparator can compare any two sets of cases for any two dimensions.

[0172] In viewing the results of the method of the present invention, when a box on a table or in a matrix or a hyperlink is selected, the case listing is generated. When a user clicks on any of the numbers, he/she is provided with a listing of each of the cases corresponding to that link. An exemplary Case List is provided in Figure 21. For each case, various information is provided, including case ID (field 2100), gender (field 2101), Manufacturer Control Code (field 2102), FDA Report Receipt Date (field 2103), Age (field 2104), Drugs (field 2105), Reactions (field 2106), Seriousness (field 2107) (Y/N or normal outcome (optional)). These columns can be sorted by clicking on their headings. If a user selects a summary view, a profile of the cases in the case list is then calculated and displayed. Additionally, if a user wishes to learn the details of a specific case, he/she can click on the case ID number of any specific case on the correlation details screen.

[0173] This Case Details screen provides detailed information on each specific case. In addition to standard information such as the patient's case ID, gender, and age, it also includes Reactions (including detailed information in the As Reported, Preferred Term, High Level Term, and High Level Group Term categories); Concomitant Drugs (each listed by Name, Dose, Route, and Suspect Status); Outcomes; Manufacturer Control Code; Manufacturer Date; Adverse Event Date; Report Type; Report Source; Case Source; and Narrative, if any. As will be appreciated, additional details can be provided. If these details are structured, all features of the invention are expandable to that dimension. If the information is unstructured, the invention can extract and structure the data using the dictionary and thesaurus facilities.

[0174] Various preferred embodiments of the invention have been described in fulfillment of the various objects of the invention. It should be recognized that these embodiments are merely illustrative of the principles of the invention. Numerous modifications and adaptations thereof will be readily apparent to those skilled in the art without departing from the spirit and scope of the present invention.

102050 985T8960